State of Hawaii Annual Summary 2014 Air Quality Data



2014 Hawaii Air Quality Data

Contents

| LIST OF TABLES | ii |
|---|-----|
| LIST OF FIGURES | iii |
| ERRATA Sheets for 2013 Data | iv |
| Section 1 INTRODUCTION | 1 |
| Section 2 DEFINITIONS | 3 |
| Section 3 SITE LOCATIONS AND DESCRIPTIONS | 7 |
| Section 4 2014 AIR QUALITY DATA | 17 |
| Section 5 2014 PM _{2.5} SPECIATION DATA | 34 |
| Section 6 AMBIENT AIR QUALITY TRENDS | 37 |

List of Tables

| Table | Title | Page |
|-------|---|------|
| 2-1 | State of Hawaii and Federal Ambient Air Quality Standards | |
| 3-1 | State of Hawaii Ambient Air Monitoring Network | |
| 3-2 | Sampling Equipment at Each Monitoring Station | 16 |
| 4-1 | 2014 Summary of the 24-Hour PM ₁₀ Averages | |
| 4-2 | Attainment Determination of the 24-Hour PM ₁₀ NAAQS | |
| 4-3 | 2014 Summary of the 24-Hour PM _{2.5} Averages: SLAMS Stations | |
| 4-4 | Attainment Determination of the 24-Hour PM _{2.5} NAAQS: SLAMS Stations | |
| 4-5 | Attainment Determination of the Annual PM _{2.5} NAAQS: SLAMS Stations | |
| 4-6 | 2014 Summary of the 24-Hour PM _{2.5} Averages: SPM Stations | |
| 4-7 | 2014 Summary of the 8-Hour O ₃ Averages | |
| 4-8 | Attainment Determination of the 8-Hour O ₃ NAAQS | |
| 4-9 | 2014 Summary of the 1-Hour and Annual NO ₂ Averages | |
| 4-10 | Attainment Determination of the 1-Hour NO ₂ NAAQS: SLAMS Stations | |
| 4-11 | 2014 Summary of the 1-Hour SO ₂ Averages NAAQS | |
| 4-12 | Attainment Determination of the 1-Hour SO ₂ NAAQS: SLAMS Stations | |
| 4-13 | 2014 Summary of the 3-Hour SO ₂ Averages | |
| 4-14 | 2014 Summary of the 24-Hour and Annual SO ₂ Averages | |
| 4-15 | 2014 Summary of the 1-Hour CO Averages | |
| 4-16 | 2014 Summary of the 8-Hour CO Averages | |
| 4-17 | 2014 Summary of the 1-Hour H ₂ S Averages (State Standard) | 26 |
| 4-18 | 2014 Summary of the Rolling 3-Month Lead Averages | |
| 4-19 | 2014 Monthly Maximum of 24-Hour PM ₁₀ Values (µg/m ³) | |
| 4-20 | 2014 Monthly Maximum of 24-Hour PM _{2.5} Values (µg/m ³) | |
| 4-21 | 2014 Monthly Maximum of 1-Hour NO ₂ Values (ppm) | |
| 4-22 | 2014 Monthly Maximum of 1-Hour CO Values (ppm) | |
| 4-23 | 2014 Monthly Maximum of 8-Hour CO Values (ppm) | |
| 4-24 | 2014 Monthly Maximum of 8-Hour O ₃ Values (ppm) | |
| 4-25 | 2014 Monthly Maximum of 1-Hour SO ₂ Values (ppm) | |
| 4-26 | 2014 Monthly Maximum of 3-Hour SO ₂ Values (ppm) | |
| 4-27 | 2014 Monthly Maximum of 24-Hour SO ₂ Values (ppm) | |
| 4-28 | 2014 Monthly Maximum of 1-Hour H ₂ S Values (ppm) | |
| 4-29 | 2014 Monthly Maximum of Rolling 3-Month Lead Values (μg/m³) | |
| 5-1 | Annual Summary of PM _{2.5} Speciation Data | |
| 5-2 | Speciation Collection and Analysis Methods | 36 |

List of Figures

| Figure | e Title | Page |
|--------|--|------|
| 3-1 | Island of Oahu Air Monitoring Stations | 7 |
| 3-2 | Island of Maui Air Monitoring Stations | |
| 3-3 | Island of Hawaii Air Monitoring Stations | 11 |
| 3-4 | Island of Kauai Air Monitoring Station | 14 |
| 6-1 | PM ₁₀ Annual Average: 2010-2014 | 38 |
| 6-2 | PM ₁₀ Maximum 24-Hour Average: 2010-2014 | 38 |
| 6-3 | PM _{2.5} Annual Average: 2010-2014 | 39 |
| 6-4 | PM _{2.5} 98 th Percentile 24-Hour Average: 2010-2014 | 39 |
| 6-5 | SO ₂ Annual Average: 2010-2014 | 40 |
| 6-6 | SO ₂ Maximum 24-Hour Average: 2010-2014 | 40 |
| 6-7 | NO ₂ Annual Average: 2010-2014 | |
| 6-8 | NO ₂ Maximum 1-Hour Average: 2010-2014 | |
| 6-9 | O ₃ Fourth Highest Daily Maximum 8-Hour Average: 2010-2014 | 42 |
| 6-10 | CO Maximum 1-Hour Average: 2010-2014 | 42 |
| 6-11 | CO Maximum 8-Hour Average: 2010-2014 | 43 |

Table 3-2 Sampling Equipment at Each Monitoring Station

| Manitanian | DM | DM | DM | - 00 | 00 | | NO | 11.0 | Land |
|-----------------------|---|--|--|---|---|---|--|---|---|
| Monitoring Station | PM ₁₀ Continuous Ambient Particulate Monitor | PM _{2.5} Manual Particulat e Monitor | PM _{2.5} Continuous Monitor | CO Continuous Gas Filter Correlation Analyzer | SO ₂ Continuous Pulsed Fluorescence Ambient Air Analyzer | O ₃ Continuous UV Photometric Analyzer | NO ₂ Continuous Chemiluminescence Analyzer | H ₂ S Continuous Pulsed Fluorescence Ambient Air Analyzer | Lead 1 in 6 Days Total Suspended Particulate Monitor |
| OAHU- Honolulu | • | | • | • | • | | | | |
| Kapolei | • | | | | | | | | • |
| Pearl City | • | | | | | | | | |
| Sand Island | | | | | | • | | | |
| MAUI- Kihei | | | | | | | | | |
| Paia | | | | | | | | | |
| HAWAII- Hilo | | | • | | • | | | | |
| Kona | | | | | | | | | |
| Mt. View | | | | | | | | | |
| Ocean View | | | | | | | | | |
| Pahala | | | | | | | | | |
| Puna E | | | | | | | | | |
| Waikoloa | | | | | | | | | |
| KAUAI- Niumalu | | | | | | | | | |

| | Maxi | mum | Annual Mean No. of 24-hour Averages Greater than 35 µg/m³ | | | | | | | | | | | | | | | |
|-------------|-------------------------|-----------------------|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 98 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | | | | | | | | | | | | | | | | | | |
| Honolulu | 17.5 | 12.4 | 5.3 ¹ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 296 | 81.1 |
| Kapolei | 16.2 | 12.0 | 2.8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 347 | 95.1 |
| Pearl City | 16.2 | 13.2 | 5.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 361 | 98.9 |
| Sand Island | 19.6 | 11.9 | 6.2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 355 | 97.3 |
| MAUI | | | | | | | | | | | | | | | | | | |
| Kihei | 25.4 | 13.1 | 5.8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 342 | 93.7 |

¹ Does not meet summary criteria, <75% data recovery in 3rd quarter

<

ERRATA Sheet for 2013 Air Data Book

The table below shows corrections (highlighted)

Table 4-6. 2013 Summary of the 24-Hour PM_{2.5} Averages: SPM Stations

| | Max | imum | Annual Mean | | No. of 24-hour Averages Greater than 35 μg/m ³ | | | | | | | | | | | | | |
|------------|-------------------------|--------------------|------------------|-----|---|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|---------------------|----------------------|---------------------|
| | 1 st High | 98 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Period s | Percent Recovery |
| HAWAII | | | | | | | | | | | | | | | | | | |
| Hilo | 22.8 | 17.1 | 7.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 358 | 98.1 |
| Kona | 28.8 | 22.6 | 13.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 356 | 97.5 |
| Mt. View | 29.8 | 16.5 | 3.2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 349 | 95.6 |
| Ocean View | 25.4 | 21.6 | 11.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 352 | 96.4 |
| Pahala | 18.4 | 14.0 | 5.4 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 357 | 97.8 |
| Waikoloa | 32.2 | 21.9 | 8.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 350 | 95.9 |
| KAUAI | | | | | | | | | | | | | | | | | | |
| Niumalu | 13.0 | 10.6 | 3.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 356 | 97.5 |
| MAUI | | | | | | | | | | | | | | | | | | |
| Paia | 18.0 | 14.3 | 7.2 ¹ | - | - | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 306 ² | 280 | 91.5 |

The special purpose stations on Hawaii island were established to monitor ambient air concentrations of PM_{2.5} from volcanic emissions. The special purpose station on Kauai was established to monitor emissions from cruise ships. The special purpose station on Kauai was established to monitor emissions from cane burning.

¹ Does not meet summary criteria, <75% data recovery in year

² Station began 3/1/2013, incomplete year

Section 1 INTRODUCTION

The Department of Health, Clean Air Branch, monitors the ambient air in the State of Hawaii for various gaseous and particulate air pollutants. The U. S. Environmental Protection Agency (EPA) has set national ambient air quality standards (NAAQS) for six criteria pollutants: carbon monoxide, nitrogen dioxide, sulfur dioxide, lead, ozone, and particulate matter (PM₁₀ and PM_{2.5}). Hawaii has also established a state ambient air standard for hydrogen sulfide. The primary purpose of the statewide monitoring network is to measure ambient air concentrations of these pollutants and ensure that these air quality standards are met. The stations are maintained and the data are collected by the Air Surveillance and Analysis Section of the State Laboratories Division.

In addition to monitoring the ambient air for criteria pollutants, the State of Hawaii also participates in the NCore multi pollutant monitoring network; the NCore station in Hawaii is located at the Kapolei air station. The NCore network addresses the following objectives:

- Timely reporting of data to public by supporting AIRNow, air quality forecasting, and other public reporting mechanisms;
- Support for development of emission strategies through air quality model evaluation and other observational methods;
- Accountability of emission strategy progress through tracking long-term trends of criteria and non-criteria pollutants and their precursors;
- Support for long-term health assessments that contribute to ongoing reviews of the NAAQS:
- Compliance through establishing nonattainment/attainment areas through comparison with the NAAQS;
- Support to scientific studies ranging across technological, health, and atmospheric process disciplines;
- Support to ecosystem assessments recognizing that national air quality networks benefit ecosystem assessments and, in turn, benefit from data specifically designed to address ecosystem analyses; and
- PM_{2.5} speciation monitoring that EPA determined to be essential for establishing a relationship between particle concentrations and adverse health effects and would provide valuable information in characterizing aerosols, determining the effectiveness of control strategies, and understanding the effects of particle pollution on atmospheric and regional haze.

Air pollution is caused by many different man-made and natural sources. There are industrial sources of pollution, such as power plants and refineries; mobile sources, such as cars, trucks, and buses; agricultural sources, such as cane burning; and natural sources, such as windblown dust and volcanic activity. In 2014, for the most part, the state maintained 14 air monitoring stations on 4 islands. Most commercial, industrial, and transportation activities and their associated air quality effects occur on Oahu, where 4 of

the stations are located. The monitoring stations on Maui are mainly to measure the air quality impacts from agricultural activities. The majority of stations are located on the island of Hawaii to measure air quality impacts from the volcano and geothermal energy production. The monitoring station on Kauai is mainly to measure the air quality impacts from cruise ships. The state's ambient air monitoring network is reviewed annually and relocations, additions and/or discontinuations can occur in the future as the need arises.

This report summarizes the validated air pollutant data collected at the 14 monitoring stations during calendar year 2014. Tabular summaries are provided which compare the measured concentrations of criteria pollutants with federal ambient air quality standards and of hydrogen sulfide with the state standard. The 2014 speciation data is also included in this report. Trend summaries of criteria pollutants parameters are shown graphically.

The Department of Health has a web site that displays near real-time air quality data updated throughout the day from the air monitoring stations. The data has not been reviewed for quality assurance and is subject to change but provides the public with viewing access to current air pollutant and meteorological information. To view this data online, go to http://health.hawaii.gov/cab and link to "Hawaii Ambient Air Quality Data."

Additionally, because emissions from the Kilauea volcano are affecting communities on the island of Hawaii on a daily basis, the Department of Health has a website dedicated to displaying short term SO₂ data from stations located on the island. It provides near real-time 15-minute SO₂ averages and advisory level guidance to help individuals protect themselves against possible health effects. To view this data online, go to www.hiso2index.info

To view this entire book as well as books from 2012 and 2013 online, go to: http://health.hawaii.gov/cab and link to "Hawaii Air Quality Data Book."

Questions or comments regarding data in this report and other air quality information should be addressed to:

Clean Air Branch Phone: (808)586-4200 Department of Health Fax: (808)586-4359

P.O. Box 3378

Honolulu, Hawaii 96801-3378

The Department of Health provides access to its programs and activities without regard to race, color, national origin (including language), age, sex, religion, or disability. Write our Affirmative Action Officer at P.O. Box 3378, Honolulu, Hawaii 96801-3378, or call (808)586-4616 (voice) within 180 days of a problem.

Cover photo is a view of part of the Ko'olau Range above 'loleka' a valley, Ko'olaupoko, Oahu.

Section 2 DEFINITIONS

98th Percentile Value The PM_{2.5} 24-hour average

The PM_{2.5} 24-hour average or the maximum daily 1-hour NO₂ average in the year below which 98% of all values fall.

99th Percentile Value

The maximum daily 1-hour SO₂ value in the year below which 99% of all values fall.

Ambient Air

The general outdoor atmosphere, external to buildings, to which the general public has access.

Ambient Air Quality Standard

A limit in the quantity and exposure to pollutants dispersed or suspended in the ambient air. Primary standards are set to protect public health, including sensitive populations such as asthmatics, children, and the elderly. Secondary standards are set to protect public welfare including protection against visibility degradation, and damage to animals, crops, vegetation and buildings.

Carbon Monoxide

Carbon monoxide (CO) is a colorless, odorless, tasteless gas under atmospheric conditions. It is produced by the incomplete combustion of carbon fuels with the majority of emissions coming from transportation sources.

CFR

Code of Federal Regulations is the codification of the general and permanent rules published in the Federal Register by the executive departments and agencies of the Federal government. Title 40 is the Protection of the Environment.

Collocated

This is a procedure required for a certain percentage of PM_{10} and $PM_{2.5}$ samplers in the monitoring network. Collocated samplers determine precision or variation in the PM_{10} or $PM_{2.5}$ concentration measurements of identical samplers run in the same location under the same sampling conditions.

Criteria Pollutants

These are the six pollutants for which the EPA has established national air quality standards. The pollutants are ozone, carbon monoxide, nitrogen dioxide, sulfur dioxide, lead and particulate matter (PM₁₀ and PM_{2.5}).

EPA

The U. S. Environmental Protection Agency; established to protect human health and the natural environment.

Hydrogen Sulfide

Hydrogen sulfide (H₂S) is a toxic, colorless gas with a characteristic "rotten egg" odor detectable at very low levels. It occurs naturally during the decomposition of organic matter, near geothermal sources and is also produced during certain industrial processes, including wastewater treatment facilities.

Micron

One micron is one millionth of a meter or approximately 1/25,000 of an inch.

 $\mu g/m^3$

Micrograms per cubic meter. This is the measurement of air quality expressed as mass per unit volume.

NAAQS

National Ambient Air Quality Standards. These are pollutant standards that the EPA has established to protect public health and welfare. NAAQS have been set for carbon monoxide, nitrogen dioxide, PM₁₀, PM_{2.5}, ozone, sulfur dioxide, and lead. These are commonly referred to as criteria pollutants.

NCore

A multi-pollutant network that integrates several advanced measurement systems for particles, pollutant gases and meteorology. Most NCore stations have been operating since the formal start of the network on January 1, 2011, including Hawaii's.

Nitrogen Dioxide

Nitrogen dioxide (NO₂) is a brownish, highly corrosive gas with a pungent odor. It is formed in the atmosphere from emissions of nitrogen oxides (NO_x). Sources of nitrogen oxides include electric utilities, industrial boilers, motor vehicle exhaust and combustion of fossil fuels. NO₂ is also a component in the atmospheric reaction that produces ground-level ozone.

Ozone

Ozone (O_3) is the main constituent in photochemical air pollution. It is formed in the atmosphere by a chemical reaction of nitrogen oxides (NO_x) and volatile organic compounds (VOC_3) in the presence of sunlight. In the upper atmosphere, O_3 shields the earth from harmful ultraviolet radiation; however, at ground level, it can cause harmful effects in humans and plants.

Particulate Matter

This refers to any solid or liquid matter dispersed in the air. Particulate matter (PM) includes dust, soot, smoke, and liquid droplets from sources such as factories, power plants, motor vehicles, construction, agricultural activities, and fires.

PM₁₀

Particulate matter that is 10 microns or less in aerodynamic diameter. These are considered "coarse" particles, generally from sources such as road and windblown dust, and crushing and grinding operations.

 $PM_{2.5}$

Particulate matter that is 2.5 microns or less in aerodynamic diameter. Considered "fine" particles, these are generally a result of fuel combustion such as from motor vehicles, utility generation and industrial facilities. Fine particles can also be formed when gases, such as sulfur dioxide and nitrogen dioxide, are chemically transformed into particles.

ppm

Parts per million is one particle in 1,000,000 other particles. It is approximately one drop in 13 gallons.

SLAMS

State and Local Air Monitoring Stations. The Clean Air Act requires that every state establish a network of air monitoring stations for criteria pollutants.

SPM

Special Purpose Monitoring stations. These are stations established to provide data for special studies in support of air program interests and activities. SPM stations supplement the SLAMS network as circumstances require and resources permit.

Sulfur Dioxide

Sulfur dioxide (SO₂) is a colorless gas that easily combines with water vapor forming sulfuric acid. Emissions of sulfur dioxide are largely from sources that burn fossil fuels such as coal and oil. In Hawaii, another major source of sulfur dioxide emissions is from the eruption of Kilauea Volcano on the Big Island.

Vog

Vog is a local term used to express volcanic smog. Vog occurs when volcanic gas and particles combine with air and sunlight to produce atmospheric haze.

Table 2-1 State and Federal Ambient Air Quality Standards

Sources: State standards HAR §11-59; Federal standards 40 CFR Part 50

| A:- | | | Standards | |
|-------------------------|---------------------|--------------------------|--|--|
| Air Pollutant | Averaging Time | Hawaii State Standard | Federal Primary Standard ^a | Federal Secondary Standard ^b |
| Carbon Monoxide | 1-hour | 9 ppm | 35 ppm | None |
| (CO) | 8-hour | 4.4 ppm | 9 ppm | None |
| Nitrogen Dioxide | 1-hour | | 0.100 ppm | |
| (NO ₂) | Annual | 0.04 ppm | 0.053 ppm | 0.053 ppm |
| PM ₁₀ | 24-hour | 150 μg/m³ | 150 μg/m³ | |
| F IVI 10 | Annual ^c | 50 μg/m³ | | |
| PM _{2.5} | 24-hour | | 35 μg/m³ | 35 μg/m³ |
| F IVI2.5 | Annual | | 12 μg/m ³ | 15 μg/m³ |
| Ozone (O ₃) | 8-hour | 0.08 ppm | 0.075 ppm | 0.075 ppm |
| | 1-hour | | 0.075 ppm | |
| Sulfur Dioxide | 3-hour | 0.5 ppm | | 0.5 ppm |
| (SO ₂) | 24-hour | 0.14 ppm | | |
| | Annual | 0.03 ppm | | |
| Lead (Pb) | Rolling 3-month | 1.5 μg/m ^{3 d} | 0.15 μg/m ³ | 0.15 μg/m ³ |
| Hydrogen Sulfide | 1-hour | 0.025 ppm | None | None |

Primary Standards set limits to protect public health, including the health of "sensitive" populations such as asthmatics, children and the elderly.

Compliance with the National Ambient Air Quality Standards

CO 1-hour: May not be exceeded more than once per year.

May not be exceeded more than once per year.

May not be exceeded more than once per year.

NO₂ 1-hour: The 3-year average of the 98th percentile daily maximum 1-hour averages must not exceed

the standard.

 NO_2 Annual: Average of all 1-hour values in the year may not exceed the level of the standard. PM_{10} 24-hour: Must not be exceeded more than one day per year, after compensating for days when

monitoring did not occur (estimated number of exceedances)

PM_{2.5} 24-hour: The 3-year average of the 98th percentile 24-hour concentrations must not exceed the level of

the standard.

PM_{2.5} Annual: The 3-year average of 24-hour values must not exceed the level of the standard.

Ozone 8-hour: The 3-year average of the fourth highest daily maximum value must not exceed the level of

the standard.

SO₂ 1-hour: The 3-year average of the 99th percentile daily maximum 1-hour averages must not exceed

the standard.

SO₂ 3-hour: Not be exceeded more than once per year. **SO₂ 24-hour:** Not be exceeded more than once per year.

SO₂ Annual: Average of all 1-hour values in the year may not exceed the level of the standard.

Lead: Average of all 24-hour values in any rolling 3-month period may not exceed the level of the

standard.

Secondary Standards set limits to protect public welfare, including protection against decreased visibility, damage to animals, crops, vegetation, and buildings.

^C Due to a lack of evidence linking health problems to long-term exposure to coarse particle pollution, EPA revoked the annual PM₁₀ standard effective December 17, 2006. However, the state still has an annual standard.

d The state standard is based on calendar quarter.

Section 3 SITE LOCATIONS AND DESCRIPTIONS

0 2.5 5 Miles
LIIIIIII

Waianae

Raneole

Campbell
Industrial Park

Honolulu

Figure 3-1: Island of Oahu Air Monitoring Stations

| Station | Name | Location | Pollutants Monitored | | | | | |
|---------------------------------|-------------------------------|--|--|--|--|--|--|--|
| 1 Honolulu 1250 Punchbowl St. I | | 1250 Punchbowl St. | PM ₁₀ , PM _{2.5} , CO, SO ₂ | | | | | |
| 2 | Sand Island | 1039 Sand Island Pkwy | O ₃ , PM _{2.5} | | | | | |
| 3 | Pearl City | 860 4th St. PM ₁₀ , PM _{2.5} | | | | | | |
| 4 | 4 Kapolei 2052 Lauwiliwili St | | PM ₁₀ , PM _{2.5} , CO, SO ₂ , NO ₂ | | | | | |
| | Kapolei NCore | 2052 Lauwiliwili St. | PM _{10-2.5} , SO _{2 trace} , NO/NO _y , CO, O ₃ , Pb, PM _{2.5} speciation, WS/WD | | | | | |

The following station descriptions include latitude and longitude in decimal degrees and altitude in meters above mean sea level.



Honolulu (DH)

| Location: | 1250 Punchbowl St., Honolulu |
|--------------|--|
| Latitude: | 21.30758 |
| Longitude: | -157.85542 |
| Altitude: | 20 m |
| Parameters: | SO ₂ , CO, PM ₁₀ , PM _{2.5} |
| Established: | February 1971 |

Brief Description:

Located in downtown Honolulu on the roof of the Department of Health building, across from the Queen's Medical Center, in a busy commercial, business and government district.



Kapolei (KA)

| Location: | 2052 Lauwiliwili St., Kapolei |
|--------------|--|
| Latitude: | 21.32374 |
| Longitude: | -158.08861 |
| Altitude: | 17.9 m |
| Parameters: | SO ₂ , CO, NO ₂ , PM ₁₀ , PM _{2.5} , |
| Parameters. | PM _{2.5} speciation, NCore |
| Established: | July 2002 |
| | · |

Brief Description:

Located in the Kapolei Business Park, southeast of the Kapolei Fire Station, next to a drainage canal that separates the park from Barber's Point.



Pearl City (PC)

| | Location: | 860 4th St., Pearl City |
|---|--------------|--------------------------------------|
| 3 | Latitude: | 21.39283 |
| ŀ | Longitude: | -157.96913 |
| i | Altitude: | 23.1 m |
| | Parameters: | PM ₁₀ , PM _{2.5} |
| , | Established: | May 1979 |
| | | |

Brief Description:

Located on the roof of the Leeward Health Center in a commercial, residential and light industrial area approximately 1.5 miles northwest of the Waiau power plant and near the Pearl Harbor Naval Complex.




Sand Island (SI)

| Location: | 1039 Sand Island Pkwy., Honolulu | | | | | | |
|--------------|------------------------------------|--|--|--|--|--|--|
| Latitude: | 21.30384 | | | | | | |
| Longitude: | -157.87712 | | | | | | |
| Altitude: | 5.3 m | | | | | | |
| Parameters: | O ₃ , PM _{2.5} | | | | | | |
| Established: | February 1981 | | | | | | |
| | | | | | | | |

Brief Description:

Located in a light industrial, commercial and recreational area approximately two miles downwind of downtown Honolulu near the entrance to the Sand Island State Recreation Area.

Figure 3-2: Island of Maui - Air Monitoring Station



| Station | Name | Location | Pollutants Monitored |
|---------|-------|-------------------|----------------------|
| 1 | Kihei | Hale Piilani Park | PM _{2.5} |
| 2 | Paia | TMK 2-5-005-05 | PM _{2.5} |



| Kihei (KH) | |
|--------------------|--------------------------|
| Location: | Hale Piilani Park, Kihei |
| Latitude: | 20.780997 |
| Longitude: | -156.44637 |
| Altitude: | 46.5 m |
| Parameters: | PM _{2.5} |
| Established: | February 1999 |
| Brief Description: | |


Located in a residential community park, next to agricultural land.



| Paia (KH) | |
|----------------------------|--------------------------|
| Location: | TMK (2)-2-5-005-05, Paia |
| Latitude: 20.902031 | |
| Longitude: | -156.370344 |
| Altitude: 80.8 m | |
| Parameters: | PM _{2.5} |
| Established: | March 2013 |
| Brief Description: | |

Located within a fenced area that contains a County of Maui water supply tank. The area is surrounded by residential and agricultural land with sugar cane fields north of the monitor.





| Station | Name | Location | Pollutants Monitored |
|---------|---------------|------------------------------|-------------------------------------|
| 1 | Hilo | 1099 Waianuenue Ave. | PM _{2.5} , SO ₂ |
| 2 | Mountain View | 17-1235 Volcano Rd. | PM _{2.5} , SO ₂ |
| 3 | Puna E | TMK (3)-1-3-28-37 (Leilani) | H ₂ S, SO ₂ |
| 4 | Pahala | 96-3150 Pikake St. | PM _{2.5} , SO ₂ |
| 5 | Ocean View | 92-6091 Orchid Mauka Circ. | PM _{2.5} , SO ₂ |
| 6 | Kona | 81-1043 Konawaena School Rd. | PM _{2.5} , SO ₂ |
| 7 | Waikoloa | TMK (3)-6-8-002-019 | PM _{2.5} , SO ₂ |



| Hilo (HL) | |
|--------------|-------------------------------------|
| Location: | 1099 Waianuenue Ave., Hilo |
| Latitude: | 19.71756 |
| Longitude: | -155.11053 |
| Altitude: | 136.8 m |
| Parameters: | SO ₂ , PM _{2.5} |
| Established: | January 1997 |
| | • |

Brief Description:

Located near the Hilo Medical Center, this station was established to monitor vog during "Kona" or southerly wind conditions.



| Kona (KN) | | |
|---|-------------------------------------|--|
| Location: 81-1043 Konawaena School | | |
| | Rd., Kona | |
| Latitude: | 19.50978 | |
| Longitude: | -155.91342 | |
| Altitude: | 517.2 m | |
| Parameters: | SO ₂ , PM _{2.5} | |
| Established: September 2005 | | |
| Brief Description: | | |

Located on the upper campus of Konawaena High School, this station monitors for vog on the west side of the island of Hawaii.



| N | Mt. View (MV) | | | | |
|--------|-----------------------|-------------------------------------|--|--|--|
| | Location: | 17-1235 Volcano Rd., Mt. View | | | |
| | Latitude: | 19.57002 | | | |
| | Longitude: | -155.08046 | | | |
| | Altitude: | 436.5 m | | | |
| | Parameters: | SO ₂ , PM _{2.5} | | | |
| | Established: | December 2010 | | | |
| To the | Brief Descript | ion: | | | |

Located on the grounds of the Mt. View Elementary School, this station was established to monitor vog during southerly wind conditions.



Ocean View (OV)

| Location: | 92-6091 Orchid Mauka Circle, | |
|--------------|-------------------------------------|--|
| | Ocean View | |
| Latitude: | 19.11756 | |
| Longitude: | -155.77814 | |
| Altitude: | 862.6 m | |
| Parameters: | SO ₂ , PM _{2.5} | |
| Established: | April 2010 | |
| | | |

Brief Description:

This station is located in Hawaii Ocean View Estates at the Ocean View fire station and monitors for volcanic emissions.



Pahala (PA)

| Location: | 96-3150 Pikake St., Pahala |
|--------------|-------------------------------------|
| Latitude: | 19.2039 |
| Longitude: | -155.48018 |
| Altitude: | 320 m |
| Parameters: | SO ₂ , PM _{2.5} |
| Established: | August 2007 |
| | |

Brief Description:

The station is on the grounds of the Kau High and Pahala Elementary School, monitoring for volcanic emissions.



Puna E (PE)

| Location: | 13-763 Leilani Ave., Pahoa | |
|--------------------|------------------------------------|--|
| Latitude: | 19.46399 | |
| Longitude: | -154.89871 | |
| Altitude: | 207.9 m | |
| Parameters: | SO ₂ , H ₂ S | |
| Established: | March 1991 | |
| Drief Description. | | |

Brief Description:

Located in the Leilani Estates residential subdivision, this station monitors for emissions from the geothermal energy facility approximately 1 mile to the northeast. The station also monitors for SO₂ emissions from the volcano during southwesterly wind conditions.




Waikoloa (WL)

| Location: | TMK 3-6-8-002-019, Waikoloa | |
|--------------|-------------------------------------|--|
| Latitude: | 19.977467 | |
| Longitude: | -155.798067 | |
| Altitude: | 180.1 m | |
| Parameters: | SO ₂ , PM _{2.5} | |
| Established: | August 2012 | |
| D : (D : (: | | |

Brief Description:

This station is located within a fenced area that contains a County of Hawaii water tank and pump house, approximately 3 km northeast of Waikoloa. This station monitors for volcanic emissions.

Figure 3-4: Island of Kauai Air Monitoring Station





| N | liumal | u (| (NI) |
|---|--------|-----|------|
| | 1 000 | 416 | n. |

| Location: | 2342 Hulemalu Road, Lihue |
|--------------|---|
| Latitude: | 21.9495 |
| Longitude: | -159.365 |
| Altitude: | 11 m |
| Parameters: | SO ₂ , CO, NO ₂ , PM _{2.5} |
| Established: | April 2011 |
| | • |

Brief Description:

Located in the Niumalu residential subdivision, this station monitors for emissions from the cruise ships in Nawiliwili Harbor approximately 1.0 mile upwind.

15

Table 3-1 State of Hawaii Ambient Air Monitoring Network

| | Po | llutar | ts Mo | nitore | ed and | Statio | on Typ | ре | | |
|--|------------------|------------------------|------------------|-----------------------|--------------------------|-----------------|------------------|--------------------|---|---|
| SITE | PM ₁₀ | PM _{2.5} | СО | O ₃ | SO ₂ | NO ₂ | H₂S | Lead | MONITORING OBJECTIVE | LOCATION SETTING |
| OAHU Honolulu Kapolei ¹ Pearl City Sand Island | S S S - | \$ \$,C \$ \$ | S S | - S - S | S S | - S - | - - - - | - S,C - - | Population Exposure Population Exposure Population Exposure Maximum Concentration (O ₃) Transport (PM _{2.5}) | Urban and Center City Suburban Urban and Center City Urban and Center City |
| MAUI Kihei Paia ² | - | S SPM | - | - | - | - | - | - | Source Impact (cane burning) Source Impact (cane burning) | Suburban Neighborhood |
| HAWAII Hilo Kona Mountain View | - | SPM SPM | - | | S S SPM | - | | - | Population Exposure Population Exposure (SO ₂)/ Maximum concentration (PM _{2.5}) Source Impact | Suburban Suburban Suburban |
| Ocean View Pahala Puna E | - | SPM SPM - SPM | - | - - - | SPM SPM SPM SPM | - - - | - SPM | - SPM | Welfare Impact (SO ₂)/ Source Impact (PM _{2.5}) Maximum concentration (SO ₂)/ Source Impact (PM _{2.5}) Source Impact (geothermal and volcano) | Rural Rural Suburban Suburban |
| Waikoloa ³ KAUAI Niumalu | - | SPM | SPM ⁴ | - | SPM | SPM | - | - | Source Impact Source Impact (cruise ships) | Suburban |

C = Collocated Site

S = (SLAMS) State and Local Air Monitoring Station

SPM = Special Purpose Monitoring Station (for monitoring vog, geothermal energy production and cruise ships)

1 Includes NCore station

Paia began operating 3/1/13
 Waikoloa station shut down 3/31/14

⁴ Niumalu CO shut down 4/25/13

Table 3-2 **Sampling Equipment at Each Monitoring Station**

| Monitoring Station | PM ₁₀ Continuous Ambient Particulate Monitor | PM _{2.5} Manual Particulate Monitor | PM _{2.5} Continuous Monitor | CO Continuous Gas Filter Correlation Analyzer | SO ₂ Continuous Pulsed Fluorescence Ambient Air Analyzer | O ₃ Continuous UV Photometric Analyzer | NO ₂ Continuous Chemiluminescence Analyzer | H₂S Continuous Pulsed Fluorescence Ambient Air Analyzer | Lead 1 in 6 Days Total Suspended Particulate Monitor |
|-----------------------|---|---|--|---|---|---|---|--|--|
| OAHU | | | | | | | | | |
| Honolulu | | | | | | | | | |
| Kapolei | • | | | • | • | • | • | | • |
| Pearl City | • | | • | | | | | | |
| Sand Island | | | • | | | • | | | |
| MAUI | | | | | | | | | |
| Kihei | | | | | | | | | |
| Paia | | | • | | | | | | |
| HAWAII | | | | | | | | | |
| Hilo | | | | | | | | | |
| Kona | | | • | | • | | | | |
| Mt. View | | | • | | • | | | | |
| Ocean View | | | | | ■. | | | | |
| Pahala | | | | | | | | | |
| Puna E | | | | | | | | | |
| Waikoloa | | | | | | | | | |
| KAUAI | | | | | | | | | |
| Niumalu | | | • | | | | • | | |

Section 4 2014 AIR QUALITY DATA

To protect the state's air quality from degradation, the Department of Health's Clean Air Branch is responsible for regulating and monitoring pollution sources to ensure that the levels of criteria pollutants remain well below the state and federal ambient air quality standards. Data collected from the ambient air network is validated by the Air Surveillance and Analysis Section to ensure that the reported data is of good quality and meets all quality control and assurance requirements.

The monitoring stations in communities near the volcano record higher levels of SO₂ and PM_{2.5}, with regular exceedances of the NAAQS for SO₂ and occasional exceedances of the NAAQS for PM_{2.5}. The EPA considers the volcano a natural, uncontrollable event and therefore the state is requesting exclusion of these NAAQS exceedances from attainment/non-attainment determination.

Excluding the exceedances due to the volcano, in 2014 the State of Hawaii was in attainment of all NAAQS.

Explanation of Summary Tables 4-1 through 4-18:

- Summaries are by pollutant and averaging period, with the number of occurrences exceeding the NAAQS or, in Table 4-17, the number of exceedances of the state H₂S standard (there is no federal H₂S standard);
- The "Maximum" is the highest and second highest valid values recorded in the year for the averaging period. For PM_{2.5}, the maximum and 98th percentile concentrations are provided and for O₃, the 4th highest daily maximum value is also displayed;
- The "Annual Mean" is the arithmetic mean of all valid values recorded in the year;
- "Possible Periods" is the total number of possible sampling periods in the year for the averaging period;
- "Valid Periods" is the total number of acceptable sampling periods after data validation;
- "Percent Recovery" represents the amount of quality data reported;
- Attainment with the NAAQS is determined according to 40 CFR 50.

Explanation of Tables 4-19 through 4-29:

- For each pollutant and averaging period, the highest concentration for each month is presented;
- The month with the highest value recorded in the year for each site is highlighted.

18

Table 4-1. 2014 Summary of the 24-Hour PM₁₀ Averages

| | Maxi | mum | Annual Mean | | N | o. of 2 | 24-hou | ır Ave | rages | Grea | ater tha | an 150 |) μg/n | ∩ ³ | | | | |
|------------|-------------------------|-------------------------|-------------------|-----|---|---------|--------|--------|-------|------|----------|--------|---------------------|------------------|---------------------|-----|------------------|------|
| | 1 st High | 2 nd High | All Hours | Jan | | | | | | | | | Possible Periods | Valid Periods | Percent Recovery | | | |
| OAHU | | | | | | | | | | | | | | | | | | |
| Honolulu | 30 | 29 | 13.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 317 | 86.8 |
| Kapolei | 32 | 32 | 15.0 ¹ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | i | - | 0 | 365 | 219 ² | 60.0 |
| Pearl City | 40 | 36 | 19.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 354 | 97.0 |

¹ Does not meet summary criteria, <75% data recovery in year

Table 4-2. Attainment Determination of the 24-Hour PM₁₀ NAAQS

| Station | Exceedances in 2012 | Exceedances in 2013 | Exceedances in 2014 | Sites in violation of the NAAQS |
|------------|---------------------|---------------------|---------------------|---------------------------------|
| Honolulu | 0 | 0 | 0 | 0 |
| Kapolei | 0 | 0 | 0 | 0 |
| Pearl City | 0 | 0 | 0 | 0 |

Attainment: The standard not to be exceeded more than once per year on average over 3 years. In 2014, Hawaii was in attainment with the 24-hour PM₁₀ NAAQS.

² No values from September to November, incomplete year

Table 4-3. 2014 Summary of the 24-Hour PM_{2.5} Averages: SLAMS Stations

| | Maxi | mum | Annual Mean | | ١ | No. of | 24-ho | ur Ave | erages | Gre | ater th | nan 35 | μg/m | 1 ³ | | | | |
|-------------|-------------------------|-----------------------|------------------|-----|-----|--------|-------|--------|--------|-----|---------|--------|------|----------------|-----|---------------------|------------------|---------------------|
| | 1 st High | 98 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | | | | | | | | | | | | | | | | | | |
| Honolulu | 12.1 | 10.2 | 4.4 ¹ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 299 | 81.9 |
| Kapolei | 14.3 | 10.7 | 3.9 ² | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | - | - | - | 0 | 365 | 202 4 | 55.3 |
| Pearl City | 15.5 | 10.3 | 3.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 342 | 93.7 |
| Sand Island | 14.1 | 10.0 | 4.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 345 | 94.5 |
| MAUI | | | | | | | | | | | | | | | | | | |
| Kihei | 14.0 | 10.4 | 3.9 ³ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 260 | 71.2 |

^{1, 2, 3,} Does not meet summary criteria, <75% data recovery in one or more quarters

Table 4-4. Attainment Determination of the 24-Hour PM_{2.5} NAAQS: SLAMS Stations

| Station | 2012 98 th value | 2013 98 th value | 2014 98 th value | 3-Year Average | Sites in violation of the NAAQS |
|-------------|-----------------------------|-----------------------------|-----------------------------|----------------|---------------------------------|
| Honolulu | 13 | 12 | 10 | 12 | 0 |
| Kapolei | 15 | 12 | 11 | 13 | 0 |
| Pearl City | 13 | 13 | 10 | 12 | 0 |
| Sand Island | 13 | 12 | 10 | 12 | 0 |
| Kihei | 14 | 13 | 10 | 12 | 0 |

Attainment: The 3-year average of the 98^{th} percentile values must be less than or equal to $35 \mu g/m^3$. In 2014, Hawaii was in attainment with the 24-hour PM_{2.5} NAAQS.

Table 4-5. Attainment Determination of the Annual PM_{2.5} NAAQS: SLAMS Stations

| Station | 2012 Ann. Avg. | 2013 Ann. Avg. | 2014 Ann. Avg. | 3-Year Average | Sites in violation of the NAAQS |
|-------------|----------------|----------------|----------------|----------------|---------------------------------|
| Honolulu | 5.1 | 5.3 | 4.4 | 4.9 | 0 |
| Kapolei | 7.1 | 2.8 | 3.9 | 4.6 | 0 |
| Pearl City | 6.3 | 5.5 | 3.9 | 5.2 | 0 |
| Sand Island | 6.2 | 6.2 | 4.5 | 5.6 | 0 |
| Kihei | 6.4 | 5.8 | 3.9 | 5.4 | 0 |

Attainment: The 3-year average of annual mean values must be less than 15 $\mu g/m^3$.

In 2014, Hawaii was in attainment with the annual PM_{2.5} NAAQS.

⁴ No values from September to November, incomplete year

Table 4-6. 2014 Summary of the 24-Hour PM_{2.5} Averages: SPM Stations

| | Maxi | mum | Annual Mean | | ١ | No. of | 24-ho | ur Ave | erages | s Gre | ater th | nan 35 | µg/m | 3 | | | | |
|------------|-------------------------|--------------------|------------------|-----|-----|--------|-------|--------|--------|-------|---------|--------|------|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 98 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| HAWAII | | | | | | | | | | | | | | | | | | |
| Hilo | 19.7 | 15.5 | 4.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 330 | 90.4 |
| Kona | 27.0 | 17.6 | 10.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 335 | 91.8 |
| Mt. View | 20.6 | 14.4 | 3.4 ¹ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 278 | 76.2 |
| Ocean View | 35.2 | 21.7 | 11.6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 315 | 86.3 |
| Pahala | 23.7 | 14.8 | 5.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 335 | 91.8 |
| Waikoloa | 42.4 | 38.8 | 9.4 | 0 | 0 | 0 | - | - | - | - | - | - | - | - | - | 90 ² | 84 | 93.3 |
| KAUAI | | | | | | | | | | | | | | | | | | |
| Niumalu | 14.5 | 10.4 | 4.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 326 | 89.3 |
| MAUI | | | | | | | | | | | | | | | | | | |
| Paia | 49.7 | 12.6 | 5.3 ³ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 299 | 81.9 |

The special purpose stations on Hawaii island were established to monitor ambient air concentrations of PM_{2.5} from volcanic emissions. The special purpose station on Kauai was established to monitor emissions from cruise ships. The special purpose station on Maui was established to monitor emissions from cane burning.

Table 4-7. 2014 Summary of the 8-Hour O₃ Averages

| | N | Maximu | m | Annual Mean | No. | No. of Daily Maximum 8-Hour Averages Greater than 0.075 ppm | | | | | | | | | | | | | |
|----------------|-------------------------|-------------------------|-------------------------|----------------|-----|---|---|---|---|---|---|---|---------------------|------------------|---------------------|---|------|-------------------|------|
| | 1 st High | 2 nd High | 4 th High | All Hours | | | | | | | | | Possible Periods | Valid Periods | Percent Recovery | | | | |
| OAHU | | | | | | | | | | | | | | | | | | | |
| Sand Island | 0.062 | 0.061 | 0.057 | 0.022 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8755 | 8081 | 92.3 |
| Kapolei | 0.050 | 0.049 | 0.046 ¹ | 0.022 | - | - | - | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8755 | 5664 ² | 64.7 |

¹ Does not meet summary criteria, <75% data recovery in year

^{1,3} Does not meet summary criteria, <75% data recovery in one or more quarters

² Station shut down 3/31/2014, incomplete year

² No values from January to March, incomplete year

Table 4-8. Attainment Determination of the 8-Hour O₃ NAAQS

| Station | 2012 4th highest | 2013 4th highest | 2014 4 th highest | 3-Year Average | Site in violation of the NAAQS |
|-------------|------------------|--------------------|------------------------------|--------------------|--------------------------------|
| Sand Island | 0.044 | 0.047 | 0.057 | 0.049 | 0 |
| Kapolei | 0.048 | 0.051 ¹ | 0.046 ¹ | 0.048 ² | 0 |

Attainment: The 3-year average of the annual 4th highest daily maximum 8-hour average must be less than or equal to 0.075 ppm. In 2014, Hawaii was in attainment with the 8-hour O₃ NAAQS.

¹ Does not meet summary criteria, <75% data recovery in year

² No values from August to December 2013 and January to March 2014, incomplete years

Table 4-9. 2014 Summary of the 1-Hour and Annual NO₂ Averages

| | | mum ·hr | Annual Mean | No. | of Da | ily Ma | ximur | m 1-Ho | our Av | /erag | es Gre | eater t | han 0 | .100 p | pm | | | |
|---------|-------------------------|--------------------|-------------|-----|-------|--------|-------|--------|--------|-------|--------|---------|-------|--------|-----|---------------------|------------------|---------------------|
| | 1 st High | 98 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAMS stations | | | | | | | | | | | | | | | | | |
| Kapolei | 0.043 | 0.025 | 0.004 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7101 | 81.1 |
| KAUAI | SPM Station | | | | | | | | | | | | | | | | | |
| Niumalu | 0.045 | 0.033 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7756 | 88.5 |

Attainment of the annual NO₂ NAAQS: The annual mean shall not exceed 0.053 ppm.

In 2014, Hawaii was in attainment with the annual NO₂ NAAQS.

Table 4-10. Attainment Determination of the 1-Hour NO2 NAAQS

| Station | 2012 98th value | 2013 98th value | 2014 98th value | 3-Year Average | Site in violation of the NAAQS |
|-----------------|-----------------|-----------------|-----------------|----------------|--------------------------------|
| Kapolei | 0.019 | 0.023 | 0.025 | 0.022 | 0 |
| Attainment. The | . 0 | - O0th | | | |

Attainment: The 3-year average of the 98th percentile values must be less than or equal to 0.100 ppm.

In 2014, Hawaii was in attainment with the 1-hour NO₂ NAAQS.

4-11. 2014 Summary of the 1-Hour SO₂ Averages

| | Maxi | mum | Annual Mean | | ١ | No. of | 1-hou | r Aver | ages | Grea | ter tha | ın 0.07 | 75 ppr | n | | | | |
|------------|-------------------------|--------------------|----------------|-----|-----|--------|-------|--------|------|------|---------|---------|--------|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 99 th % | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAMS | Stations | i | | | | | | | | | | | | | | | |
| Honolulu | 0.018 | 0.006 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7801 | 89.1 |
| Kapolei | 0.024 | 0.021 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 6702 | 76.5 |
| HAWAII | SPM St | ations (s | ee NOTE) | | | | | | | | | | | | | | | |
| Hilo | 0.331 | 0.144 | 0.004 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 4 | 8760 | 7897 | 90.1 |
| Kona | 0.051 | 0.035 | 0.003 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7930 | 90.5 |
| Mt. View | 0.691 | 0.312 | 0.005 | 8 | 6 | 2 | 0 | 0 | 0 | 0 | 1 | 1 | 2 | 2 | 5 | 8760 | 8052 | 91.9 |
| Ocean View | 0.866 | 0.379 | 0.013 | 12 | 8 | 8 | 9 | 8 | 7 | 7 | 4 | 3 | 8 | 10 | 9 | 8760 | 7858 | 89.7 |
| Pahala | 0.645 | 0.444 | 0.024 | 17 | 9 | 14 | 25 | 13 | 16 | 18 | 8 | 12 | 11 | 12 | 19 | 8760 | 8259 | 94.3 |
| Puna E | 0.106 | 0.034 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 8760 | 8102 | 92.5 |
| Waikoloa | 0.020 | 0.020 | 0.001 | 0 | 0 | 0 | - | - | - | - | - | - | - | - | - | 2160 ¹ | 2122 | 98.2 |
| KAUAI | SPM St | ation | | | | | | | | | | | | | | | | |
| Niumalu | 0.052 | 0.036 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7663 | 87.5 |

Attainment: The 3-year average of the 99th percentile values must be less than or equal to 0.075 ppm. Effective June 2, 2010. In 2014, Hawaii was in attainment with the 1-hour SO₂ NAAQS (SLAMS stations only).

NOTE: The SPM stations on Hawaii Island were established to monitor ambient air concentrations of SO_2 from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 1-hour NAAQS from attainment determinations. The SPM station on Kauai was established to monitor emissions from cruise ships.

¹ Station shut down 3/31/14, incomplete year

23

Table 4-12. Attainment Determination of the 1-Hour SO₂ NAAQS: SLAMS Stations

| | 2012 99th value | 2013 99th value | 2014 99th value | 3-Year Average | Violation of the NAAQS |
|--------------------------------------|--------------------|-----------------|--------------------|----------------|------------------------|
| OAHU SLAMS stations | | | | | N= NO Y= YES |
| Honolulu | 0.016 | 0.007 | 0.006 | 0.009 | N |
| Kapolei | 0.007 | 0.009 | 0.021 1 | 0.012 | N |
| HAWAII SPM stations (SEE note) | | | | | |
| Hilo | 0.227 | 0.169 | 0.144 ¹ | 0.180 | Υ |
| Kona | 0.064 | 0.042 | 0.035 | 0.047 | N |
| Mt. View | 0.277 | 0.248 | 0.312 | 0.279 | Υ |
| Ocean View | 0.754 | 0.466 | 0.379 | 0.533 | Υ |
| Pahala | 0.935 | 0.757 | 0.444 | 0.712 | Υ |
| Puna E | 0.008 | 0.016 | 0.034 | 0.019 | N |
| Waikoloa | 0.032 ² | 0.020 | 0.020 ³ | 0.024 | N |
| KAUAI SPM station | | | | | |
| Niumalu | 0.041 1 | 0.025 | 0.036 | 0.034 | N N |

Attainment: The 3-year average of the 99th percentile values must be less than or equal to 0.075 ppm. Effective June 2, 2010. In 2014, Hawaii was in attainment with the 1-hour SO₂ NAAQS (SLAMS stations only).

NOTE: The SPM stations on Hawaii Island were established to monitor ambient air concentrations of SO_2 from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 1-hour NAAQS from attainment determinations. The SPM station on Kauai was established to monitor emissions from cruise ships.

¹ Does not meet summary criteria, < 75% data recovery in one or more quarters ³ Station shut down 3/31/14, incomplete year ² Station began operations 7/1/12, incomplete year

Table 4-13. 2014 Summary of the 3-Hour SO₂ Averages

| | Maxi | mum | Annual Mean | | | No. o | f 3-ho | ur Ave | erages | Gre | ater th | an 0.5 | 5 ppm | | | | | |
|------------|-------------------------|-------------------------|----------------|-----|-----|-------|--------|--------|--------|-----|---------|--------|-------|-----|-----|---------------------|-------------------|---------------------|
| | 1 st High | 2 nd High | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAMS | stations | 5 | | | | | | | | | | | | | | | |
| Honolulu | 0.015 | 0.014 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2512 | 86.0 |
| Kapolei | 0.019 | 0.013 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2142 ¹ | 73.4 |
| HAWAII | SPM st | ations (s | ee NOTE) | | | | | | | | | | | | | | | |
| Hilo | 0.285 | 0.248 | 0.004 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2529 | 86.6 |
| Kona | 0.036 | 0.036 | 0.003 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2538 | 86.9 |
| Mt. View | 0.276 | 0.261 | 0.005 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2601 | 89.1 |
| Ocean View | 0.389 | 0.272 | 0.013 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2506 | 85.8 |
| Pahala | 0.458 | 0.277 | 0.024 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2673 | 91.5 |
| Puna E | 0.058 | 0.047 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2571 | 88.0 |
| Waikoloa | 0.012 | 0.011 | 0.001 | 0 | 0 | 0 | - | 1 | - | - | - | - | - | ı | - | 720 ² | 688 | 95.6 |
| KAUAI | SPM st | ation | | | | | | | | | | | | | | | | |
| Niumalu | 0.046 | 0.039 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2920 | 2478 | 84.9 |

Attainment: 3-hour values not to exceed 0.5 ppm more than once per year.

In 2014, Hawaii was in attainment with the 3-hour SO₂ NAAQS (SLAMS stations only).

NOTE: The SPM stations on Hawaii island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 3-hour NAAQS from attainment determinations.

¹ Does not meet summary criteria, < 75% data recovery in one or more quarters

² Station shut down 3/31/14, incomplete year

Table 4-14. 2014 Summary of the 24-Hour and Annual SO₂ Averages

| | Max | imum | Annual Mean | | N | o. of 2 | 24-hou | ır Ave | rages | Grea | iter tha | an 0.1 | 40 pp | m | | | | |
|------------|-------------------------|-------------------------|----------------|-----|-----|---------|--------|--------|-------|------|----------|--------|-------|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 2 nd High | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAM | S Statio | ns | | | | | | | | | | | | | | | _ |
| Honolulu | 0.008 | 0.006 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 327 | 89.6 |
| Kapolei | 0.008 | 0.005 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 273 ¹ | 74.8 |
| HAWAII | SPM S | Stations | (see NOTE) | | | | | | | | | | | | | | | |
| Hilo | 0.077 | 0.057 | 0.004 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 332 | 91.0 |
| Kona | 0.015 | 0.011 | 0.003 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 333 | 91.2 |
| Mt. View | 0.079 | 0.070 | 0.005 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 339 | 92.9 |
| Ocean View | 0.090 | 0.086 | 0.013 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 331 | 90.7 |
| Pahala | 0.126 | 0.102 | 0.024 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 347 | 95.1 |
| Puna E | 0.011 | 0.009 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 342 | 93.7 |
| Waikoloa | 0.007 | 0.007 | 0.001 | 0 | 0 | 0 | - | - | - | - | - | - | - | ı | - | 90 ² | 90 | 100 |
| KAUAI | SPM S | Station | | | | | | | | | | | | | | | | |
| Niumalu | 0.019 | 0.016 | 0.002 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 365 | 323 | 88.5 |

Attainment: 24-hour values not to exceed 0.14 ppm more than once per year.

In 2014, Hawaii was in attainment of the state 24-hour SO₂ standard (SLAMS stations only).

NOTE: The SPM stations on Hawaii island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 24-hour NAAQS from attainment determinations.

Attainment: Annual average (from SLAMS stations only) not to exceed 0.03 ppm.

In 2014, Hawaii was in attainment of the state annual SO₂ standard.

NOTE: The SPM stations on Hawaii island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the annual NAAQS from attainment determinations.

¹ Does not meet summary criteria, <75% data recovery in year

^{2,} Station shut down 3/31/14, incomplete year

Table 4-15. 2014 Summary of the 1-Hour CO Averages

| | Maxi | mum | Annual Mean | | | No. of | 1-ho | ur Ave | rages | Grea | ater th | an 35 | ppm | | | | | |
|----------|-------------------------|-------------------------|----------------|-----|-----|--------|------|--------|-------|------|---------|-------|-----|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 2 nd High | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAMS | S station | s | | | | | | | | | | | | | | | |
| Honolulu | 2.0 | 1.8 | 0.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7626 | 87.1 |
| Kapolei | 2.7 | 1.4 | 0.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7528 | 85.9 |

Attainment: 1-hour values not to exceed 35 ppm more than once per year.

In 2014, Hawaii was in attainment with the 1-hour CO NAAQS.

Table 4-16. 2014 Summary of the 8-Hour CO Averages

| | | | 1 4510 | | | | | . , | | | <u> </u> | | J. J. | , | | | | |
|----------|-------------------------|-------------------------|----------------|-----|-----|-------|--------|------------|--------|-------|----------|---------|-------|-----|-----|---------------------|------------------|---------------------|
| | Maxi | mum | Annual Mean | | | No. o | f 8-ho | ur Ave | erages | s Gre | ater th | nan 9 p | opm | | | | | |
| | 1 st High | 2 nd High | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| OAHU | SLAMS | S station | s | | | | | | | | | | | | | | | |
| Honolulu | 1.0 | 1.0 | 0.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8755 | 7675 | 87.7 |
| Kapolei | 1.3 | 1.3 | 0.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8755 | 7486 | 85.5 |

Attainment: 8-hour values not to exceed 9 ppm more than once per year.

In 2014, Hawaii was in attainment with the 8-hour CO NAAQS.

Table 4-17. 2014 Summary of the 1-Hour H₂S Averages (State Standard)

| | Maxi | mum | Annual Mean | | | No. o | f 1-ho | ur Aver | ages (| Great | er thar | า 0.02 | 5 ppm | | | | | |
|--------|-------------------------|-------------------------|-------------|-----|-----|-------|--------|---------|--------|-------|---------|--------|-------|-----|-----|---------------------|------------------|---------------------|
| | 1 st High | 2 nd High | All Hours | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Possible Periods | Valid Periods | Percent Recovery |
| HAWAII | | | | | | | | | | | | | | | | | | |
| Puna E | 0.006 | 0.005 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 8760 | 7905 | 90.2 |

Attainment of the state standard: 1-hour values not to exceed 0.025 ppm.

In 2014, Hawaii was in attainment of the state 1-hour H₂S standard.

Table 4-18. 2014 Summary of the Rolling 3-Month Lead Averages

| | Maxi | mum | Annual Mean | | ı | No. of | 3-Mor | nth Ave | erages | Grea | ter tha | n 0.15 | μg/m ³ | 3 | | | | |
|---------|-------|-------------------------|----------------|-----|---|--------|-------|---------|--------|------|---------|--------|-------------------|---|-----|---------------------|------------------|---------------------|
| | 1 st | ond | | | Rolling 3-Month period ending in the month of | | | | | | | | | | | D 11-1 - | \ | D |
| | High | 2 nd High | All Hours | Jan | <u> </u> | | | | | | | | | | Dec | Possible Periods | Valid Periods | Percent Recovery |
| HAWAII | | | | | | | | | | | | | | | | | | |
| Kapolei | 0.001 | 0.001 | 0.001 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 61 | 49 | 80.3 |

Attainment: Maximum 3-month average concentration for a 3-year period must be less than or equal to 0.15 µg/m³.

Note: Sampling for lead conducted 1 in 6 days. Sampling began 1/1/2012.

Table 4-19. 2014 Monthly Maximum of 24-Hour PM₁₀ Values (μg/m³)

The month with the highest value in the year is highlighted

The state and federal 24-hr PM₁₀ standard is 150 μ g/m³

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Honolulu | 29 | 19 | 25 | 30 | 27 | 18 | 16 | 18 | 17 | 17 | 22 | 23 |
| Kapolei ¹ | 32 | 30 | 30 | 32 | 24 | 28 | 14 | 21 | - | - | - | 24 |
| Pearl City | 36 | 30 | 40 | 35 | 33 | 30 | 18 | 23 | 27 | 28 | 33 | 34 |

¹ Kapolei station no values from September to November, incomplete year

Table 4-20. 2014 Monthly Maximum of 24-Hour PM_{2.5} Values (μg/m³)

The month with the highest value in the year is highlighted

The federal 24-hr PM_{2.5} standard is $35 \mu g/m^3$

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| SLAMS Stations | | | | | | | | | | | | |
| Honolulu | 12 | 11 | 6 | 12 | 12 | 6 | 6 | 9 | 8 | 9 | 8 | 10 |
| Kapolei ¹ | 10 | 9 | 11 | 14 | 9 | 7 | 6 | 8 | - | - | - | 7 |
| Pearl City | 16 | 13 | 8 | 12 | 12 | 6 | 5 | 7 | 5 | 7 | 6 | 11 |
| Sand Island | 12 | 11 | 13 | 10 | 11 | 6 | 5 | 7 | 5 | 9 | 8 | 14 |
| Kihei | 8 | 10 | 9 | 11 | 11 | 9 | 6 | 6 | 3 | 8 | 6 | 14 |
| SPM Stations | | | | | | | | | | | | |
| Niumalu (cruise ships) | 7 | 8 | 9 | 15 | 11 | 9 | 7 | 5 | 10 | 10 | 11 | 12 |
| Hilo (volcano) | 20 | 12 | 15 | 8 | 7 | 6 | 7 | 6 | 5 | 7 | 16 | 17 |
| Kona (volcano) | 18 | 18 | 27 | 16 | 14 | 16 | 16 | 16 | 16 | 15 | 16 | 21 |
| Mt. View (volcano) | 21 | 11 | 17 | 10 | 7 | 6 | 4 | 5 | 6 | 12 | 11 | 10 |
| Ocean View (volcano) | 16 | 18 | 35 | 18 | 18 | 17 | 16 | 15 | 18 | 22 | 24 | 23 |
| Pahala (volcano) | 24 | 11 | 20 | 13 | 11 | 8 | 8 | 8 | 8 | 5 | 14 | 19 |
| Waikoloa 1 (volcano) | 15 | 20 | 42 | = | - | - | - | - | - | - | - | - |
| Paia (cane burning) | 13 | 24 | 11 | 12 | 12 | 7 | 5 | 12 | 50 | 7 | 13 | 11 |

¹ Kapolei station no values from September to November, incomplete year

² Waikoloa station shut down 3/31/14

Table 4-21. 2014 Monthly Maximum of 1-Hour NO₂ Values (ppm)

The month with the highest value in the year is highlighted

The federal 1-hour standard for NO₂ is 0.100 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Kapolei | 0.029 | 0.029 | 0.028 | 0.020 | 0.025 | 0.010 | 0.015 | 0.043 | 0.018 | 0.016 | 0.023 | 0.030 |
| Niumalu | 0.029 | 0.029 | 0.026 | 0.045 | 0.033 | 0.028 | 0.022 | 0.022 | 0.025 | 0.022 | 0.037 | 0.036 |

Table 4-22. 2014 Monthly Maximum of 1-Hour CO Values (ppm)

The month with the highest value in the year is highlighted

The federal 1-hr CO standard is 35 ppm, the state standard is 9ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Honolulu | 1.0 | 1.2 | 2.0 | 0.9 | 1.3 | 0.6 | 0.8 | 0.4 | 0.8 | 0.9 | 1.3 | 1.1 |
| Kapolei | 1.2 | 1.0 | 1.4 | 1.1 | 1.0 | 1.3 | 0.4 | 0.5 | 1.1 | 0.7 | 2.7 | 1.0 |

Table 4-23. 2014 Monthly Maximum of 8-Hour CO Values (ppm)

The month with the highest value in the year is highlighted

The federal 8-hr CO standard is 9 ppm, the state standard is 4.4 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Honolulu | 0.8 | 0.8 | 0.9 | 0.8 | 1.0 | 0.5 | 0.6 | 0.3 | 0.6 | 0.6 | 0.6 | 0.8 |
| Kapolei | 1.0 | 1.0 | 1.1 | 1.0 | 0.9 | 1.3 | 0.3 | 0.4 | 0.6 | 0.6 | 0.9 | 0.7 |

4-24. 2014 Monthly Maximum of 8-Hour O₃ Values (ppm)

The month with the highest value in the year is highlighted

The federal 8-hr O₃ standard is 0.075 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|----------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Sand Island | 0.047 | 0.049 | 0.057 | 0.062 | 0.048 | 0.037 | 0.030 | 0.028 | 0.033 | 0.036 | 0.043 | 0.045 |
| Kapolei NCore ¹ | - | - | = | 0.050 | 0.048 | 0.036 | 0.028 | 0.032 | 0.031 | 0.032 | 0.047 | 0.044 |

¹ Kapolei station no values from January to March, incomplete year

Table 4-25. 2014 Monthly Maximum of 1-Hour SO₂ Values (ppm)

The month with the highest value in the year is highlighted

The federal 1-hr SO₂ standard is 0.075 ppm (75 ppb)

| | | | | | | | | | | - 1-1 (| | |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|---------|-------|-------|
| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
| SLAMS Stations | | | | | | | | | | | | |
| Honolulu | 0.018 | 0.006 | 0.004 | 0.004 | 0.004 | 0.002 | 0.005 | 0.002 | 0.002 | 0.001 | 0.002 | 0.002 |
| Kapolei | 0.023 | 0.010 | 0.008 | 0.008 | 0.004 | 0.005 | 0.020 | 0.005 | 0.005 | 0.024 | 0.012 | 0.009 |
| SPM Stations (see NOTE) | | | | | | | | | | | | |
| Niumalu (cruise ships) | 0.009 | 0.011 | 0.026 | 0.050 | 0.017 | 0.032 | 0.031 | 0.021 | 0.016 | 0.016 | 0.035 | 0.052 |
| Hilo (volcano) | 0.075 | 0.305 | 0.058 | 0.029 | 0.051 | 0.017 | 0.009 | 0.045 | 0.032 | 0.062 | 0.087 | 0.331 |
| Kona (volcano) | 0.024 | 0.051 | 0.045 | 0.019 | 0.015 | 0.020 | 0.020 | 0.014 | 0.009 | 0.011 | 0.029 | 0.018 |
| Mt. View (volcano) | 0.354 | 0.196 | 0.691 | 0.023 | 0.074 | 0.008 | 0.038 | 0.101 | 0.093 | 0.168 | 0.101 | 0.362 |
| Ocean View (volcano) | 0.866 | 0.439 | 0.107 | 0.137 | 0.716 | 0.232 | 0.159 | 0.154 | 0.238 | 0.202 | 0.223 | 0.211 |
| Pahala (volcano) | 0.451 | 0.399 | 0.645 | 0.444 | 0.271 | 0.330 | 0.345 | 0.224 | 0.204 | 0.381 | 0.538 | 0.392 |
| Puna E (volcano) | 0.043 | 0.010 | 0.010 | 0.004 | 0.003 | 0.001 | 0.001 | 0.007 | 0.001 | 0.001 | 0.075 | 0.106 |
| Waikoloa 1 (volcano) | 0.014 | 0.020 | 0.020 | - | - | - | - | - | - | - | - | - |

NOTE: The SPM stations on Hawaii Island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Although Hilo and Kona stations are designated SLAMS, the values are still mostly attributed to volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 1-hour NAAQS from attainment determinations.

¹ Waikoloa station shut down 3/31/14

Table 4-26. 2014 Monthly Maximum of 3-Hour SO₂ Values (ppm)

The month with the highest value in the year is highlighted

The state and federal 3-hr SO₂ standard is 0.5 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| SLAMS Stations | | | | | | | | | | | | |
| Honolulu | 0.015 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.003 | 0.002 | 0.002 | 0.000 | 0.001 | 0.001 |
| Kapolei | 0.019 | 0.005 | 0.005 | 0.005 | 0.004 | 0.005 | 0.005 | 0.002 | 0.003 | 0.012 | 0.005 | 0.007 |
| SPM Stations (see NOTE) | | | | | | | | | | | | |
| Niumalu (cruise ships) | 0.007 | 0.006 | 0.015 | 0.046 | 0.014 | 0.022 | 0.027 | 0.016 | 0.011 | 0.013 | 0.031 | 0.039 |
| Hilo (volcano) | 0.052 | 0.285 | 0.050 | 0.020 | 0.032 | 0.009 | 0.005 | 0.020 | 0.011 | 0.026 | 0.079 | 0.248 |
| Kona (volcano) | 0.020 | 0.036 | 0.036 | 0.017 | 0.011 | 0.017 | 0.017 | 0.011 | 0.007 | 0.008 | 0.018 | 0.014 |
| Mt. View (volcano) | 0.160 | 0.097 | 0.276 | 0.018 | 0.048 | 0.005 | 0.015 | 0.089 | 0.034 | 0.074 | 0.092 | 0.261 |
| Ocean View (volcano) | 0.389 | 0.236 | 0.087 | 0.090 | 0.272 | 0.168 | 0.085 | 0.123 | 0.176 | 0.113 | 0.182 | 0.135 |
| Pahala (volcano) | 0.268 | 0.252 | 0.458 | 0.277 | 0.183 | 0.257 | 0.270 | 0.172 | 0.139 | 0.166 | 0.227 | 0.274 |
| Puna E (volcano) | 0.032 | 0.009 | 0.007 | 0.003 | 0.002 | 0.001 | 0.001 | 0.006 | 0.001 | 0.001 | 0.047 | 0.058 |
| Waikoloa 1 (volcano) | 0.011 | 0.012 | 0.010 | - | - | - | - | - | - | - | - | - |

NOTE: The SPM stations on Hawaii Island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 3-hour NAAQS from attainment determinations.

¹ Waikoloa station shut down 3/31/14

Table 4-27. 2014 Monthly Maximum of 24-Hour SO₂ Values (ppm)

The month with the highest value in the year is highlighted

The state and federal 24-hr SO₂ standard is 0.14 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| SLAMS Stations | | | | | | | | | | | | |
| Honolulu | 0.008 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.000 | 0.001 | 0.001 |
| Kapolei | 0.008 | 0.003 | 0.003 | 0.002 | 0.004 | 0.004 | 0.003 | 0.000 | 0.001 | 0.004 | 0.003 | 0.003 |
| SPM Stations (see NOTE) | | | | | | | | | | | | |
| Niumalu (cruise ships) | 0.002 | 0.002 | 0.003 | 0.019 | 0.006 | 0.010 | 0.012 | 0.007 | 0.004 | 0.005 | 0.016 | 0.007 |
| Hilo (volcano) | 0.024 | 0.057 | 0.015 | 0.006 | 0.008 | 0.003 | 0.002 | 0.005 | 0.004 | 0.009 | 0.028 | 0.077 |
| Kona (volcano) | 0.011 | 0.015 | 0.009 | 0.006 | 0.005 | 0.008 | 0.006 | 0.005 | 0.004 | 0.004 | 0.009 | 0.010 |
| Mt. View (volcano) | 0.079 | 0.022 | 0.068 | 0.007 | 0.012 | 0.003 | 0.006 | 0.012 | 0.008 | 0.020 | 0.031 | 0.070 |
| Ocean View (volcano) | 0.090 | 0.076 | 0.036 | 0.037 | 0.053 | 0.059 | 0.024 | 0.027 | 0.030 | 0.030 | 0.086 | 0.054 |
| Pahala (volcano) | 0.102 | 0.065 | 0.126 | 0.093 | 0.055 | 0.059 | 0.081 | 0.062 | 0.044 | 0.069 | 0.098 | 0.089 |
| Puna E (volcano) | 0.009 | 0.004 | 0.003 | 0.002 | 0.002 | 0.001 | 0.001 | 0.001 | 0.000 | 0.000 | 0.009 | 0.011 |
| Waikoloa 1 (volcano) | 0.007 | 0.007 | 0.003 | - | - | - | - | - | - | - | - | - |

NOTE: The SPM stations on Hawaii Island were established to monitor ambient air concentrations of SO₂ from volcanic emissions. Volcanic eruptions are considered natural events and therefore EPA may exclude the exceedances of the 24-hour NAAQS from attainment determinations.

¹ Waikoloa station shut down 3/31/14

Table 4-28. 2014 Monthly Maximum of 1-Hour H₂S Values (ppm)

The month with the highest value in the year is highlighted

The state 1-hour H₂S standard is .025 ppm

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Puna E | 0.001 | 0.002 | 0.006 | 0.002 | 0.002 | 0.001 | 0.003 | 0.003 | 0.003 | 0.004 | 0.002 | 0.005 |

Table 4-29. 2014 Monthly Maximum of Rolling 3-Month Lead Values (μg/m³)
The month with the highest value in the year is highlighted

The federal rolling 3-month lead standard is 0.15 μg/m³

| Station | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Kapolei NCore (1 in 6 days) | 0.002 | 0.002 | 0.001 | 0.002 | 0.002 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.003 |

Section 5 2014 PM_{2.5} SPECIATION DATA

Atmospheric aerosols are solid or liquid particles suspended in air that come directly from a variety of sources (primary) or are formed by chemical reactions (secondary). Primary and secondary particles tend to have long lifetimes in the atmosphere and can travel long distances, up to hundreds or perhaps thousands of miles. Sources include dust from roads, construction, and agriculture; combustion particles from motor vehicles, electric utilities and agricultural burning; and particles from natural sources such as the ocean or volcano.

Most of the PM_{2.5} is a combination of the following components: sulfates, nitrates, ammonium, elemental carbon, organic compounds, water and metals. The EPA selected target particulates of interest based on data use objectives, primary constituents of PM_{2.5}, and the capability and availability of current analytical methods.

The filter-based speciation sampler collects samples once every 3 days for analyses performed by an EPA contract laboratory. The speciation sampler is located at the Kapolei NCore monitoring station.

Table 5-1 lists the parameters measured, highest and second highest values recorded in the year, the annual arithmetic mean of all valid samples and the total number of samples collected in the year. Table 5-2 lists the analysis methods for each parameter.

With the exception of lead, there are no ambient air quality standards for the individual components of speciated PM_{2.5}.

For more information on EPA's speciation program, go to: www.epa.gov/ttn/amtic/speciepg.html

Table 5-1. Annual Summary of PM_{2.5} Speciation Data

| Parameter | 1 st High (µg/m³) | 2 nd High (μg/m³) | Annual Mean (µg/m³) | No. of Samples | Percent Recovery |
|------------------|---------------------------------|---------------------------------|------------------------|-------------------|---------------------|
| CARBON | | | | | |
| Organic Carbon | 1.120 | 0.949 | 0.4085 | 98 | 81 |
| Elemental Carbon | 0.351 | 0.312 | 0.1174 | 98 | 81 |
| METALS | | | | | |
| Aluminum | 0.150 | 0.116 | 0.0219 | 98 | 81 |
| Antimony | 0.092 | 0.036 | 0.0193 | 98 | 81 |
| Arsenic | 0.001 | 0.001 | 0.0002 | 98 | 81 |
| Barium | 0.030 | 0.030 | 0.0058 | 98 | 81 |
| Bromine | 0.005 | 0.004 | 0.0014 | 98 | 81 |
| Cadmium | 0.022 | 0.012 | 0.0009 | 98 | 81 |
| Calcium | 0.118 | 0.113 | 0.0295 | 98 | 81 |
| Cerium | 0.044 | 0.043 | 0.0048 | 98 | 81 |
| Cesium | 0.023 | 0.023 | 0.0086 | 98 | 81 |
| Chlorine | 1.590 | 1.370 | 0.4531 | 98 | 81 |
| Chromium | 0.029 | 0.016 | 0.0013 | 98 | 81 |
| Cobalt | 0.002 | 0.001 | 0.0007 | 98 | 81 |
| Copper | 0.011 | 0.006 | 0.0015 | 98 | 81 |
| Indium | 0.018 | 0.017 | 0.093 | 98 | 81 |
| Iron | 0.133 | 0.106 | 0.0287 | 98 | 81 |
| Lead | 0.004 | 0.004 | 0.0003 | 98 | 81 |
| Magnesium | 0.138 | 0.138 | 0.0384 | 98 | 81 |
| Manganese | 0.002 | 0.002 | 0.0003 | 98 | 81 |
| Nickel | 0.017 | 0.011 | 0.0026 | 98 | 81 |
| Phosphorus | 0.008 | 0.008 | 0.0057 | 98 | 81 |
| Potassium | 0.542 | 0.084 | 0.0286 | 98 | 81 |
| Rubidium | 0.002 | 0.002 | 0.009 | 98 | 81 |
| Selenium | 0.001 | 0.001 | 0.0011 | 98 | 81 |
| Silicon | 0.329 | 0.254 | 0.0461 | 98 | 81 |
| Silver | 0.019 | 0.019 | 0.0069 | 98 | 81 |
| Sodium | 1.040 | 0.912 | 0.3129 | 98 | 81 |
| Strontium | 0.011 | 0.003 | 0.0013 | 98 | 81 |
| Sulfur | 2.110 | 1.440 | 0.3107 | 98 | 81 |
| Tin | 0.050 | 0.037 | 0.0132 | 98 | 81 |
| Titanium | 0.014 | 0.011 | 0.0027 | 98 | 81 |
| Vanadium | 0.011 | 0.009 | 0.0021 | 98 | 81 |
| Zinc | 0.012 | 0.008 | 0.0016 | 98 | 81 |
| Zirconium | 0.012 | 0.012 | 0.0029 | 98 | 81 |

Table 5-1 Continued

| Parameter | 1 st High (µg/m³) | 2 nd High (µg/m³) | Annual Mean (µg/m³) | No. of Samples | Percent Recovery |
|---------------|---------------------------------|---------------------------------|------------------------|-------------------|---------------------|
| IONS | | | | | |
| Ammonium Ion | 1.25 | 0.82 | 0.096 | 97 | 81 |
| Potassium Ion | 0.51 | 0.06 | 0.021 | 97 | 81 |
| Sodium Ion | 1.17 | 0.94 | 0.367 | 97 | 81 |
| Total Nitrate | 0.97 | 0.45 | 0.187 | 97 | 81 |
| Sulfate | 6.57 | 4.47 | 0.959 | 97 | 81 |

 Table 5-2.
 Speciation Collection and Analysis Methods

| Parameter | Collection Method | Analysis Method |
|-----------|----------------------------|--------------------------------------|
| Carbon | URG 300N Quartz Filter | Thermal Optical Transmittance |
| Metals | Met-One SASS Teflon Filter | Energy Dispersive X-Ray Fluorescence |
| lons | Met-One SASS Nylon Filter | Ion Chromatography |

Section 6 AMBIENT AIR QUALITY TRENDS

The following graphs illustrate 5-year trends for PM₁₀, PM_{2.5}, SO₂, NO₂, O₃, and CO from 2010 to 2014 at all SLAMS stations monitoring for those pollutants.

Figures 6-1 and 6-2 are graphs of the PM₁₀ annual and maximum 24-hour averages.

Figure 6-3 is the graph of the PM_{2.5} annual averages. Attainment of the PM_{2.5} 24-hour standard is based on the 98th percentile value at each station, which is depicted in Figure 6-4.


Figures 6-5 and 6-6 are graphs of the SO₂ annual and maximum 24-hour averages.


Figure 6-7 and 6-8 shows the annual and maximum 1-hour averages of NO₂ compared to the federal NAAQS.


Attainment of the 8-hour ozone standard is achieved by averaging 3 years of the fourth highest daily maximum 8-hour average concentrations, which must not exceed 0.075 ppm (standard effective May 27, 2008). Figure 6-9 is a graph of the fourth highest daily maximum values recorded at the Sand Island and Kapolei (since 2011) ozone monitoring stations in the past five years.


The graphs for 1-hour and 8-hour carbon monoxide (figures 6-10 and 6-11, respectively) represent the maximum 1-hour or 8-hour values recorded in the year.


Criteria pollutant levels remain below state and federal ambient air quality standards at all SLAMS stations in the state.

















Figure 6-6. SO₂ Maximum 24-Hour Average: 2010-2014




Figure 6-7. NO₂ Annual Average: 2010-2014




Figure 6-8. NO₂ Maximum 1-Hour Average: 2010-2014

